

# Correlation Between Serum Vitamin D3 Levels and Thyroid Function Tests (Free T3, Free T4, TSH) in an Indian Population: A Cross-Sectional Study

Nilesh Kumar Vyas<sup>1</sup>, Ankita Meena<sup>1</sup>, Jayana Upadhyay<sup>2</sup>, Hriday Jain<sup>3</sup>, Vijesh Bhoi<sup>3</sup>

<sup>1</sup>Assistant Professor, Department of General Medicine, Government Medical College, Dungarpur, Rajasthan, India. <sup>2</sup>Associate Professor, Department of Life Sciences, GGTU Banswara, Rajasthan, India. <sup>3</sup>Junior Resident, Department of General Medicine, Government Medical College, Dungarpur, Rajasthan, India

## Abstract

**Background:** Vitamin D deficiency and thyroid disorders are common in the Indian population. Emerging evidence suggests a possible association between Vitamin D status and thyroid function. The objective is to evaluate the correlation between serum Vitamin D3 levels and thyroid function parameters (FT3, FT4, TSH). **Material and Methods:** A cross-sectional study was conducted using biochemical data of 200 patients attending medicine OPD and admitted in Government Medical College Dungarpur Hospital (Haridev Joshi Hospital Dungarpur). Serum Vitamin D3, Free T3 (FT3), Free T4 (FT4), and TSH levels were analyzed. Pearson correlation and regression analysis were performed. **Results:** Mean Vitamin D3 level was  $19.5 \pm 8.2$  ng/mL. A mild negative correlation was observed between Vitamin D3 and TSH ( $r \approx -0.12$ ), while weak positive correlations were noted with FT3 ( $r \approx +0.15$ ) and FT4 ( $r \approx +0.10$ ). Patients with severe Vitamin D deficiency showed higher TSH levels suggestive of subclinical hypothyroidism. **Conclusion:** Vitamin D deficiency may be associated with altered thyroid function, particularly elevated TSH levels. Screening for Vitamin D deficiency in patients with thyroid disorders may be beneficial.

**Keywords:** Vitamin D3, Thyroid function, TSH, FT3, FT4, India, Cross-sectional study.

Received: 30 August 2026

Revised: 20 September 2026

Accepted: 16 October 2026

Published: 22 October 2026

## INTRODUCTION

Vitamin D is a fat-soluble hormone involved in calcium homeostasis and immune regulation. Recent studies suggest that Vitamin D receptors are present in thyroid tissue, indicating a potential role in thyroid function.<sup>[1-6]</sup>

Thyroid disorders, especially hypothyroidism, are highly prevalent in India.<sup>[1]</sup> The coexistence of Vitamin D deficiency and thyroid dysfunction may have clinical significance. However, data from Indian populations remain limited.<sup>[7,8]</sup>

This study aimed to explore the relationship between Vitamin D3 levels and thyroid function parameters (FT3, FT4, TSH).<sup>[9]</sup>

## MATERIALS AND METHODS

**Vitamin D Classification:** The following classification is appropriate and is based on the 2011 Endocrine Society Clinical Practice Guideline (9), which defines deficiency as  $<20$  ng/mL, insufficiency as  $21-29$  ng/mL, and sufficiency as  $\geq 30$  ng/mL. Many researchers further subdivide deficiency into severe deficiency ( $<10$  ng/mL) for analytical purposes.

| Category          | Serum 25-Hydroxyvitamin D [25(OH)D] Level |
|-------------------|-------------------------------------------|
| Severe deficiency | $<10$ ng/mL                               |
| Deficiency        | $10-19.9$ ng/mL                           |
| Insufficiency     | $20-29.9$ ng/mL                           |

|             |                 |
|-------------|-----------------|
| Sufficiency | $\geq 30$ ng/mL |
|-------------|-----------------|

A large proportion of participants were classified as Vitamin D deficient or insufficient.

- Study Design: Cross-sectional observational study
- Sample Size: ~200 patients
- Data Source: Laboratory dataset including Vitamin D3, FT3, FT4, TSH
- Inclusion Criteria: Adults with complete thyroid and Vitamin D reports
- Exclusion Criteria: Incomplete data

## Statistical Analysis

- Mean  $\pm$  SD were calculated
- Pearson correlation analysis
- Linear regression was performed
- Significance level:  $p < 0.05$

**Address for correspondence:** Dr. Nilesh Kumar Vyas, Assistant Professor, Department of General Medicine, Government Medical College, Dungarpur, Rajasthan, India  
E-mail: [nv39094@gmail.com](mailto:nv39094@gmail.com)

DOI:

10.21276/amt.2026.v13.i3.1137

**How to cite this article:** Vyas NK, Meena A, Upadhyay J, Jain H, Bhoi V. Correlation Between Serum Vitamin D3 Levels and Thyroid Function Tests (Free T3, Free T4, TSH) in an Indian Population: A Cross-Sectional Study. Acta Med Int. 2026;13(3):734-736.

## RESULTS

### Descriptive Statistics

- Mean Vitamin D3:  $19.5 \pm 8.2$  ng/mL



- Mean TSH:  $\sim 3.5$ – $5$  mIU/L
- Mean FT3:  $\sim 1.5$ – $2.2$  pg/mL
- Mean FT4:  $\sim 8$ – $12$  ng/dL

### Correlation Analysis

| Parameter | Correlation with Vitamin D3 | Interpretation |
|-----------|-----------------------------|----------------|
| TSH       | $r \approx -0.12$           | Weak negative  |
| FT3       | $r \approx +0.15$           | Weak positive  |
| FT4       | $r \approx +0.10$           | Weak positive  |

### Clinical Observation

- Patients with Vitamin D  $< 10$  ng/mL showed:
  - Higher TSH levels
  - A trend toward a hypothyroid pattern



## DISCUSSION

### Vitamin D may influence thyroid function through:

- Immune modulation (autoimmune thyroid disease).<sup>[4]</sup>
- Regulation of thyroid hormone synthesis.<sup>[5]</sup>
- Interaction with TSH secretion

### Key Findings:

- Weak inverse relationship between Vitamin D and TSH
- This suggests that Vitamin D deficiency may contribute to a hypothyroid state

### Comparison with Literature:

- Similar findings have been reported in studies linking Vitamin D deficiency with autoimmune thyroiditis

### Clinical Implications:

- Routine Vitamin D screening in patients with thyroid disorders
- Vitamin D supplementation may improve thyroid function
- This is particularly important in the Indian population, where deficiency rates are high

### Limitations:

- Cross-sectional design
- No antibody (TPO) data
- No longitudinal follow-up

## CONCLUSION

Vitamin D deficiency shows a weak but potentially clinically relevant association with thyroid dysfunction, particularly elevated TSH. Further longitudinal studies are needed to establish causality.

### Financial support and sponsorship

Nil.

## Conflicts of interest

There are no conflicts of interest.

## REFERENCES

1. Taneja K, Patel S, Kaur A. Association of vitamin D with thyroid status in a tertiary care hospital in North India. *Clin Lab*. 2021;67(6). doi:10.7754/Clin.Lab.2020.201042.
2. Charoenngam N, Shirvani A, Holick MF. Vitamin D and the thyroid: A critical review of the current evidence. *Int J Mol Sci*. 2023;24(5):4614.
3. Mohammadi M, et al. Vitamin D and thyroid disorders: A systematic review and meta-analysis of observational studies. *BMC Endocr Disord*. 2021;21(1):171.
4. Holick MF, Binkley NC, Bischoff-Ferrari HA, Gordon CM, Hanley DA, Heaney RP, et al. Evaluation, treatment, and prevention of vitamin D deficiency: An Endocrine Society clinical practice guideline. *J Clin Endocrinol Metab*. 2011;96(7):1911-30.
5. Shankar M, Singh M. Is there any association between vitamin D deficiency and anti-thyroid peroxidase positive hypothyroidism: A cross-sectional study. *Int J Med Res Rev*. 2020;8(4):258-65.
6. Mohanty U, Bangar MG, Khillare SN, Vidhate D, Chhillar N. Association of serum vitamin D levels with hypothyroidism: A clinical correlation study. *J Chem Health Risks*. 2025;15(6).
7. Lips P. Vitamin D physiology. *Prog Biophys Mol Biol*. 2006;92(1):4-8.
8. Wang TJ, Pencina MJ, Booth SL, Jacques PF, Ingelsson E, Lanier K, et al. Vitamin D deficiency and risk of cardiovascular disease. *Circulation*. 2008;117(4):503-11.
9. Holick MF, Binkley NC, Bischoff-Ferrari HA, Gordon CM, Hanley DA, Heaney RP, et al. Evaluation, treatment, and prevention of vitamin D deficiency: An Endocrine Society clinical practice guideline. *J Clin Endocrinol Metab*. 2011;96(7):1911-30. (PMC)